

Internet Electronic Journal of **Molecular Design**

June 2005, Volume 4, Number 6, Pages 393–412

Editor: Ovidiu Ivanciuc

Proceedings of the Internet Electronic Conference of Molecular Design 2004
IECMD 2004, November 29 – December 12, 2004

QSAR Study on 5-*N*-Substituted-2-(Substituted Benzenesulphonyl) Glutamines as Antitumor Agents through Synthesis and Biological Evaluation: Part III

Bikash Debnath, Soma Samanta, Shovanlal Gayen, Anindya Basu, Balaram Ghosh,
and Tarun Jha

Division of Pharmaceutical and Medicinal Chemistry, Department of Pharmaceutical Technology,
P.O. Box 17020, Jadavpur University, Kolkata–700032, India

Received: November 24, 2004; Revised: March 9, 2005; Accepted: March 23, 2005; Published: June 30, 2005

Citation of the article:

B. Debnath, S. Samanta, S. Gayen, A. Basu, B. Ghosh, and T. Jha, QSAR Study on 5-*N*-Substituted-2-(Substituted Benzenesulphonyl) Glutamines as Antitumor Agents through Synthesis and Biological Evaluation: Part III, *Internet Electron. J. Mol. Des.* **2005**, *4*, 393–412, <http://www.biochempress.com>.

QSAR Study on 5-*N*-Substituted-2-(Substituted Benzenesulphonyl) Glutamines as Antitumor Agents through Synthesis and Biological Evaluation: Part III[#]

Bikash Debnath, Soma Samanta, Shovanlal Gayen, Anindya Basu, Balaram Ghosh, and Tarun Jha *

Division of Pharmaceutical and Medicinal Chemistry, Department of Pharmaceutical Technology, P.O. Box 17020, Jadavpur University, Kolkata-700032, India

Received: November 24, 2004; Revised: March 9, 2005; Accepted: March 23, 2005; Published: June 30, 2005

Internet Electron. J. Mol. Des. 2005, 4 (6), 393–412

Abstract

Motivation. The importance of the non-essential amino acid, glutamine, to the proliferation of human tumour cells was well established. It is one of the two major, if not the only, substrate of cancer. It helps in tumor cell growth by supplying its amide nitrogen atom in the biosynthesis of other amino acids, purine, and pyrimidine bases. Tumor is a “nitrogen trap” as well as “glutamine trap”. Hence, the efforts were made to synthesize series of glutamine analogs, evaluate these biologically and QSAR study was performed to explore the substitutional requirements essential for improved anticancer activity.

Method. QSAR study was performed using Log of percentage tumor weight inhibition as dependent parameter and physicochemical parameter, ETSA indices and indicator variable as independent parameters through multiple linear regression analysis.

Results. Some of the compounds showed promising anticancer activity. This study resulted some QSAR models with 86.49, 83.61, 88.52, 88.41 and 88.00% of explained variances. All these models showed more than 80% predicted variance. QSAR study revealed that aliphatic substitution of the glutamine analogs might have played an important role in the hydrophobic/dispersive interaction with the possible glutamine receptor. This study also showed that field effect at R₁ position and resonance effect at R₂ position might increase anticancer activity. Some of the atoms of the general structure were identified as pharmacophore. At least one free hydrogen in amide moiety of glutamine analogs might be essential for the anticancer activity.

Conclusions. This study throws some light in the structural requirements essential for improved anticancer activity and will help to find out substituents for future synthesis of this type of analogs.

Keywords. Glutamine; anticancer agents; synthesis; screening; quantitative structure–activity relationships; physicochemical parameters; ETSA indices.

Abbreviations and notations

QSAR, quantitative structure–activity relationships

ETSA, Electrotopological State Atom

[#] Presented in part at the Internet Electronic Conference of Molecular Design 2004, IECMD 2004.

* Correspondence author; phone: +91-33-2414 6677/+91-33-2453 0945; fax: +91-33-2414 6677/+91-33-2414 6927; E-mail: tjjpharm@yahoo.com; drtarunjha@yahoo.com.

1 INTRODUCTION

Neoplastic transformation is accompanied by adaptive increases in nucleotide and protein synthesis. The high rates of protein synthesis in rapidly growing tumors require a continuous supply of both essential and nonessential amino acids [1]. It was observed that tumors assimilate not only the nitrogen from the diet, but also the nitrogen from host proteins, raising the concept of tumors as “nitrogen traps”, actively competing with the host for nitrogen compounds [1]. As glutamine is the most abundant amino acid in the body and the main vehicle for circulation of ammonia in a nontoxic form [2], it is considered that tumors behave indeed as “glutamine traps” [3,4]. After glutamine gains access to the cytoplasm, it must be transported into mitochondria. Where it is acted upon by glutaminase, an enzyme requiring high phosphate concentrations to be fully active. The high concentrations of inorganic phosphate found in the mitochondria of tumor cells could explain the high activity of tumor glutaminase *in vivo* [1]. In fact, experimental evidence supports the correlation of glutaminase activity with the extent of malignant proliferation [2,3]. The importance of the non-essential amino acid, glutamine, to the proliferation of human tumor cells was well established [3,5,6]. It is one of the two major, if not the only, substrate of cancer. The other substrate is glucose, the only circulating sugar, which is essential for the growth of normal and neoplastic cells. The presence of a tumor produces great changes in host glutamine metabolism in such a way that host nitrogen metabolism is accommodated to the tumor-enhanced requirements of glutamine [7–10]. Glutamine is also essential for the culture of many cell types. All of the cells studied had a high activity of phosphate-dependent glutaminase and were found to utilize glutamine from the culture medium during long-term culture. The rate of cell proliferation, determined by [6–3H]-thymidine incorporation, was dependent on glutamine concentration [1,11]. Considering the importance of this amino acid in cancer it has prompted us to explore the glutamine analogs for their possible anticancer activity.

In continuation of our previously reported work [12–17] on synthesis, biological evaluation and quantitative structure–activity relationship (QSAR) studies on some derivatives and analogs of glutamine as possible anticancer agents, 32 new 5-*N*-substituted-2-(substituted benzenesulphonyl) glutamines, as shown in Figure 1, were synthesized. These compounds were biologically evaluated for anticancer activity. QSAR studies, using percentage inhibition of tumor weight considered as the biological activity parameter, were performed on all thirty-two glutamine analogs. The study was done to explore the substitutional requirements essential for the improved anticancer activity.





Figure 1. General structure of 5-*N*-substituted-2-(substituted benzenesulphonyl)–glutamines 5–36.

2 MATERIALS AND METHODS

2.1 Synthesis

Thirty-two 5-*N*-substituted-2-(substituted benzenesulphonyl) glutamines were synthesized. All the reagents used for the synthesis were of AR grade and commercially available from SD Fine Chemicals, Fluka, Sigma Aldrich, Rankem.



2.1.1 Chemistry

Synthesis of 5-*N*-substituted-2-(substituted benzenesulphonyl)-glutamine analogs were carried out according to Scheme 1. Synthesis was started with chlorosulphonylation [18] of substituted benzenes **1** to get corresponding sulphonyl chlorides **2**. This sulphonyl halide **2** proved to be a versatile synthon [13–15,17] in the subsequent step in the preparation of substituted benzenesulphonyl glutamic acids **3**. With the application of Schotten–Bauman reaction [19], 2-(substituted benzenesulphonyl) glutamic acids **3** were prepared by one-step condensation of with L-glutamic acid. In this reaction alkaline medium was maintained to remove the hydrochloric acid, which was formed during condensation. Reaction of the resulting intermediates **3** with acetyl chloride afforded cyclized acid intermediates 1-(substituted benzenesulphonyl)-5-oxopyrrolidine-2-carboxylic acids **4**. Aminolysis [20] of the cyclized acid **4** with various amines afforded the corresponding glutamines **5–36**.

2.1.2 General synthetic procedure

2.1.2.1. Method 1. Substituted benzenesulphonyl chloride (2a–2d). To a mixture of substituted benzene (**1a–1d**: 0.1 mole) in chloroform (50 ml) in a 500 ml flask equipped with a dropping funnel, a thermometer and reflux condenser, chlorosulphonic acid (0.25 mole) was added dropwise over a period of 45 min to 60 min. The reaction mixture was mechanically stirred at 0 °C in a bath containing freezing mixture of ice and salt. Chlorosulphonic acid was added in such a rate that the temperature of the reaction mixture does not exceed 5 °C. In case of 2,5-dichloro benzenesulphonyl chloride (**2b**), the temperature was maintained at 90–100 °C. After the complete addition of chlorosulphonic acid, the reaction mixture was stirred for another 45 min at room temperature and the mixture was poured on to crushed ice. The product was extracted with three 50-ml portions of chloroform, dried overnight over anhydrous sodium sulphate. Chloroform was distilled off. The product was sufficiently pure which was not attempted for further purification. It had been taken for the next step.

2.1.2.2. Method 2. 2-(Substituted benzenesulphonyl) glutamic acid (3a–3d). L-glutamic acid (0.1 mole) was taken in a 250-ml conical flask and sodium hydroxide solution (2N) was added slowly till glutamic acid dissolved and the mixture become distinctly alkaline to phenolphthalein. The reaction mixture was stirred on a mechanical stirrer and the temperature was maintained at 70°C using hot water-bath. Substituted benzenesulphonyl chloride (0.11 moles) was added in small portions with constant stirring and sodium hydroxide (2N) was added time-to-time to keep the reaction mixture alkaline. The reaction was continued until a clear homogeneous solution resulted and thin layer chromatography showed the reaction was complete. After the reaction was over, it was allowed to cool to room temperature and filtered to separate undissolved solid matter, if any.

The filtrate was acidified with concentrated hydrochloric acid and saturated with sodium chloride. The product was extracted with three 50 ml portions of ethyl acetate. Ethyl acetate layer was washed with brine solution (15 ml) and dried overnight over anhydrous sodium sulphate. The solvent was distilled off to get the desired diacid (**3a–3d**).

2.1.2.3. Method 3. 1-(Substituted benzenesulphonyl)-5-oxopyrrolidine-2-carboxylic acid (4a–4d). 2-(Substituted benzenesulphonyl) glutamic acid (**3a–3d**: 0.01 mole) was taken in 100 ml round bottomed flask, fitted with reflux condenser and calcium chloride guard tube. Acetyl chloride (0.025 mole) was added to it and refluxed for 2 hr on boiling water bath. The completion of the reaction was tested by thin layer chromatography. After the reaction was completed, the reaction mixture was cooled to room temperature and poured onto crushed ice with continuous stirring. The precipitated product was filtered and recrystallized from water with charcoal treatment.

2.1.2.4. Method 4. 5-N-Substituted 2-(substituted benzenesulphonyl) glutamines (5–36). In a 50 ml of loosely stoppered conical flask, 1-(substituted benzene sulphonyl)-5-oxopyrrolidine-2-carboxylic acid (**4a–4d**: 0.01 mole) was suspended in 20 ml of water. To this, excess of amines (0.025 mole) were added and allowed to stand overnight. The reaction mixture was concentrated over steam bath to remove excess of amines. It was cooled to room temperature and chilled in an ice bath. The mixture was acidified with 6N hydrochloric acid. The precipitate was filtered and the residue was washed with cold water and finally recrystallized from dilute ethanol with charcoal treatment.

2.1.3 Characterization

Melting points of all the compounds were measured on a capillary melting point apparatus and were uncorrected. All the compounds were characterized qualitatively and quantitatively by performing both analytical and spectrophotometric analysis. The infrared spectra were recorded on BUCK M500 quick scanning Infrared spectrophotometer using KBr pellets. Running the spectrum of 0.05mm polystyrene film did the finer calibration of the machine. The frequencies were expressed in cm^{-1} . Proton Nuclear Magnetic Resonance (^1H NMR) spectra were collected at 25°C in the pulsed Fourier Transformation mode on Bruker DRX 300 MHz spectrophotometers using the solvents described and was consistent with the proposed structures. Chemical shifts are reported in δ ppm (parts per million) relative to Tetramethyl Silane for deuterated dimethylsulfoxide ($\text{DMSO}-d_6$). Signals are quoted as *s* (singlet), *d* (doublet), *t* (triplet) and *m* (multiplet). The mass spectra (FAB) were recorded on JEOL–JMS–SX–102. PNBA (*p*-nitrobenzyl alcohol) was used as matrix (M^+), which showed $\text{M}+1$ peak at 154, $2\text{M}+1$ peak at 307. Elemental or microanalyses (C, H, N) of the compounds was performed on 2400 Series II CHN analyzer of Perkin–Elmer. Reactions were monitor by analytical thin layer chromatography performed on silica gel G plates. The spots were

located keeping the TLC plates in iodine chamber. Physical data of the intermediate compounds and final compounds are summarized in Table 1 and 2 respectively. Mass-, IR-, and Proton NMR spectroscopic as well as microanalyses data of the final compounds are shown in Table 3.

Table 1. Physical data of intermediate compounds

Cpd ^a	R ₁	R ₂	R ₃	M.P.(°C)	%Yield	Molecular Formula	MW
2a	H	Br	H	73–75	77.90	C ₆ H ₄ O ₂ SClBr	255.52
2b	Cl	H	Cl	35–37	80.00	C ₆ H ₃ O ₂ SCl ₃	245.51
2c	CH ₃	CH ₃	H	29–31	70.52	C ₈ H ₉ O ₂ SCl	204.68
2d	H	<i>t</i> -C ₄ H ₉	H	56–58	68.64	C ₁₀ H ₁₃ O ₂ SCl	232.73
3a	H	Br	H	148–150	80.84	C ₁₁ H ₁₂ NO ₆ SBr	366.19
3b	Cl	H	Cl	188–190	68.56	C ₁₁ H ₁₁ NO ₆ SCl ₂	356.18
3c	CH ₃	CH ₃	H	153–155	72.52	C ₁₃ H ₁₇ NO ₆ S	315.34
3d	H	<i>t</i> -C ₄ H ₉	H	160–164	57.25	C ₁₅ H ₂₁ NO ₆ S	343.40
4a	H	Br	H	95–97	64.00	C ₁₁ H ₁₀ NO ₅ SBr	348.18
4b	Cl	H	Cl	184–186	74.00	C ₁₁ H ₉ NO ₅ SCl ₂	338.16
4c	CH ₃	CH ₃	H	180–182	48.65	C ₁₃ H ₁₅ NO ₅ S	297.33
4d	H	<i>t</i> -C ₄ H ₉	H	139–141	62.24	C ₁₅ H ₁₉ NO ₅ S	325.38

^a Compound number

Table 2. Physical data of 5-*N*-substituted-2-(substituted benzenesulphonyl)-glutamines 5–36

Cpd	R ₁	R ₂	R ₃	R ₄	R ₅	M.P. (°C)	%Yield	Molecular formula	M.W.
5	H	Br	H	<i>i</i> -C ₃ H ₇	H	193–195	64.32	C ₁₄ H ₁₉ N ₂ O ₅ SBr	407.28
6	H	Br	H	<i>c</i> -C ₆ H ₁₁	H	227–229	56.87	C ₁₇ H ₂₃ N ₂ O ₅ SBr	447.35
7	H	Br	H	C ₆ H ₅ CH ₂	H	198–200	77.42	C ₁₈ H ₁₉ N ₂ O ₅ SBr	455.32
8	H	Br	H	CH ₃	CH ₃	135–137	58.24	C ₁₃ H ₁₇ N ₂ O ₅ SBr	393.26
9	Cl	H	Cl	CH ₃	H	162–164	84.36	C ₁₂ H ₁₄ N ₂ O ₅ SCl ₂	369.22
10	Cl	H	Cl	<i>n</i> -C ₃ H ₇	H	152–154	68.54	C ₁₄ H ₁₈ N ₂ O ₅ SCl ₂	397.28
11	Cl	H	Cl	<i>i</i> -C ₃ H ₇	H	195–197	52.14	C ₁₄ H ₁₈ N ₂ O ₅ SCl ₂	397.28
12	Cl	H	Cl	<i>n</i> -C ₄ H ₉	H	119–121	84.78	C ₁₅ H ₂₀ N ₂ O ₅ SCl ₂	411.30
13	Cl	H	Cl	<i>i</i> -C ₄ H ₉	H	156–158	56.34	C ₁₅ H ₂₀ N ₂ O ₅ SCl ₂	411.30
14	Cl	H	Cl	<i>n</i> -C ₆ H ₁₃	H	140–142	74.52	C ₁₇ H ₂₄ N ₂ O ₅ SCl ₂	439.35
15	Cl	H	Cl	<i>c</i> -C ₆ H ₁₁	H	164–166	67.35	C ₁₇ H ₂₂ N ₂ O ₅ SCl ₂	437.34
16	Cl	H	Cl	C ₆ H ₅	H	148–150	65.32	C ₁₇ H ₁₆ N ₂ O ₅ SCl ₂	431.29
17	Cl	H	Cl	C ₆ H ₅ CH ₂	H	151–153	58.32	C ₁₈ H ₁₈ N ₂ O ₅ SCl ₂	445.32
18	Cl	H	Cl	CH ₃	CH ₃	118–120	48.32	C ₁₃ H ₁₆ N ₂ O ₅ SCl ₂	383.25
19	Cl	H	Cl	<i>i</i> -C ₃ H ₇	<i>i</i> -C ₃ H ₇	68–70	44.64	C ₁₇ H ₂₄ N ₂ O ₅ SCl ₂	439.35
20	CH ₃	CH ₃	H	H	H	142–144	78.84	C ₁₃ H ₁₈ N ₂ O ₅ S	314.36
21	CH ₃	CH ₃	H	<i>n</i> -C ₄ H ₉	H	120–122	84.56	C ₁₇ H ₂₆ N ₂ O ₅ S	370.47
22	CH ₃	CH ₃	H	C ₂ H ₅	C ₂ H ₅	153–155	53.32	C ₁₇ H ₂₆ N ₂ O ₅ S	370.47
23	H	<i>t</i> -C ₄ H ₉	H	H	H	132–134	68.38	C ₁₅ H ₂₂ N ₂ O ₅ S	342.41
24	H	<i>t</i> -C ₄ H ₉	H	CH ₃	H	155–157	56.64	C ₁₆ H ₂₄ N ₂ O ₅ S	356.44
25	H	<i>t</i> -C ₄ H ₉	H	C ₂ H ₅	H	107–109	58.36	C ₁₇ H ₂₆ N ₂ O ₅ S	370.47
26	H	<i>t</i> -C ₄ H ₉	H	<i>n</i> -C ₃ H ₇	H	134–136	75.54	C ₁₈ H ₂₈ N ₂ O ₅ S	384.49
27	H	<i>t</i> -C ₄ H ₉	H	<i>i</i> -C ₃ H ₇	H	134–136	72.25	C ₁₈ H ₂₈ N ₂ O ₅ S	384.49
28	H	<i>t</i> -C ₄ H ₉	H	<i>n</i> -C ₄ H ₉	H	151–153	89.94	C ₁₉ H ₃₀ N ₂ O ₅ S	398.52
29	H	<i>t</i> -C ₄ H ₉	H	<i>i</i> -C ₄ H ₉	H	143–145	76.64	C ₁₉ H ₃₀ N ₂ O ₅ S	398.52
30	H	<i>t</i> -C ₄ H ₉	H	<i>n</i> -C ₆ H ₁₃	H	143–145	84.32	C ₂₁ H ₃₄ N ₂ O ₅ S	426.57
31	H	<i>t</i> -C ₄ H ₉	H	<i>c</i> -C ₆ H ₁₁	H	170–172	88.34	C ₂₁ H ₃₂ N ₂ O ₅ S	424.56
32	H	<i>t</i> -C ₄ H ₉	H	C ₆ H ₅	H	184–186	74.45	C ₂₁ H ₂₆ N ₂ O ₅ S	418.51
33	H	<i>t</i> -C ₄ H ₉	H	C ₆ H ₅ CH ₂	H	108–110	90.92	C ₂₂ H ₂₈ N ₂ O ₅ S	432.53
34	H	<i>t</i> -C ₄ H ₉	H	CH ₃	CH ₃	103–105	52.24	C ₁₇ H ₂₆ N ₂ O ₅ S	370.47
35	H	<i>t</i> -C ₄ H ₉	H	C ₂ H ₅	C ₂ H ₅	82–84	48.95	C ₁₉ H ₃₀ N ₂ O ₅ S	398.52
36	H	<i>t</i> -C ₄ H ₉	H	<i>i</i> -C ₃ H ₇	<i>i</i> -C ₃ H ₇	130–132	52.36	C ₂₁ H ₃₄ N ₂ O ₅ S	426.57

Table 3. Mass, IR and Proton NMR spectroscopic as well as CHN analysis data of the final compounds (**5–36**)

Cpd	Mass (FAB)	IR (KBr, cm ⁻¹)	¹ HNMR (300 MHz, DMSO- <i>d</i> ₆)	C,H,N: %calcd/found		
				C	H	N
5	M + H ⁺ peak at m/z 408	3318, 3114 (N–H str of CONH), 3022 (Ar–C–H str), 2872 (ali C–H str), 1706 (C=O str), 1562, 1442 (ali C–H def), 1332 & 1160 (S=O str of SO ₂ NH), 973, 790 & 744 (Ar–C–H def)	12.66 (s, 1H, COOH), 8.46 (d, 1H, SO ₂ NH), 8.28–7.98 (m, 4H, H–2', H–3', H–5', H–6'), 7.74 (m, 1H, CONH), 3.68 (m, 1H, H–2), 2.92 (m, 1H, N–CH–1''), 2.00 (m, 2H, H ₂ –4), 1.86 (m, 1H, H _A –3), 1.68 (m, 1H, H _B –3), 1.12–0.96 (m, 6H, CH ₃ –2'', CH ₃ –3'')	41.29 41.20	4.70 4.68	6.88 6.71
6	M + H ⁺ peak at m/z 448	3314, 3104 (N–H str of CONH), 3028 (Ar–C–H str), 2874 (ali C–H str), 1705 (C=O str), 1564, 1445 (ali C–H def), 1336 & 1162 (S=O str of SO ₂ NH), 967, 794 & 748 (Ar–C–H def)	12.74 (s, 1H, COOH), 8.40 (d, 1H, SO ₂ NH), 8.12–7.88 (m, 4H, H–2', H–3', H–5', H–6'), 7.58 (m, 1H, CONH), 3.72 (m, 1H, H–2), 2.10 (m, 2H, H ₂ –4), 1.82 (m, 1H, H _A –3), 1.62 (m, 1H, H _B –3), 1.42–1.12 (m, 11H, cyclohexyl protons)	45.64 45.60	5.18 5.08	6.26 6.18
7	M + H ⁺ peak at m/z 456	3322, 3112 (N–H str of CONH), 3032 (Ar–C–H str), 2868 (ali C–H str), 1694 (C=O str), 1560, 1444 (ali C–H def), 1332 & 1160 (S=O str of SO ₂ NH), 975, 798 & 756 (Ar–C–H def)	12.72 (s, 1H, COOH), 8.48 (d, 1H, SO ₂ NH), 8.20–7.94 (m, 4H, H–2', H–3', H–5', H–6'), 7.78 (m, 1H, CONH), 7.64–7.48 (5H, Phenyl protons), 4.24 (m, 2H, CH ₂ –Ph), 3.78 (m, 1H, H–2), 2.10 (m, 2H, H ₂ –4), 1.86 (m, 1H, H _A –3), 1.70 (m, 1H, H _B –3)	47.48 47.52	4.21 4.26	6.15 6.12
8	M + H ⁺ peak at m/z 394	3028 (Ar–C–H str), 2836 (ali C–H str), 1706 (C=O str), 1565, 1448 (ali C–H def), 1326 & 1160 (S=O str of SO ₂ NH), 978, 796 & 750 (Ar–C–H def)	12.64 (s, 1H, COOH), 8.52 (d, 1H, SO ₂ NH), 8.22–7.96 (m, 4H, H–2', H–3', H–5', H–6'), 3.74 (m, 1H, H–2), 3.18–3.00 (m, 6H, CH ₃ –1'', CH ₃ –2''), 2.08 (m, 2H, H ₂ –4), 1.88 (m, 1H, H _A –3), 1.62 (m, 1H, H _B –3)	39.70 39.58	4.36 4.32	7.12 7.06
9	M + H ⁺ peak at m/z 370	3310, 3108 (N–H str of CONH), 3032 (Ar–C–H str), 2882 (ali C–H str), 1695 (C=O str), 1562, 1442 (ali C–H def), 1336 & 1164 (S=O str of SO ₂ NH), 974, 796 & 752 (Ar–C–H def)	12.68 (s, 1H, COOH), 8.52 (d, 1H, SO ₂ NH), 8.26–8.16 (m, 2H, H–3', H–6'), 7.85 (m, 1H, H–4'), 7.78 (m, 1H, CONH), 3.74m, 1H, H–2), 3.02 (m, 3H, CH ₃ –1''), 2.02 (m, 2H, H ₂ –4), 1.84 (m, 1H, H _A –3), 1.65 (m, 1H, H _B –3)	39.04 38.96	3.82 3.76	7.59 7.52
10	M + H ⁺ peak at m/z 398	3320, 3118 (N–H str of CONH), 3024 (Ar–C–H str), 2882 (ali C–H str), 1700 (C=O str), 1552, 1440 (ali C–H def), 1338 & 1164 (S=O str of SO ₂ NH), 975, 798 & 752 (Ar–C–H def)	12.72 (s, 1H, COOH), 8.40 (d, 1H, SO ₂ NH), 8.12–8.04 (m, 2H, H–3', H–6'), 7.74 (m, 1H, H–4'), 7.60 (m, 1H, CONH), 3.76 (m, 1H, H–2), 3.00 (m, 2H, N–CH ₂ –1''), 2.18 (m, 2H, H ₂ –4), 1.88 (m, 1H, H _A –3), 1.68 (m, 1H, H _B –3), 1.32 (m, 2H, CH ₂ –2''), 0.90 (m, 3H, CH ₃ –3'')	42.33 42.18	4.57 4.52	7.05 6.98
11	M + H ⁺ peak at m/z 398	3312, 3100 (N–H str of CONH), 3022 (Ar–C–H str), 2876 (ali C–H str), 1695 (C=O str), 1562, 1436 (ali C–H def), 1334 & 1162 (S=O str of SO ₂ NH), 976, 790 & 746 (Ar–C–H def)	12.70 (s, 1H, COOH), 8.44 (d, 1H, SO ₂ NH), 8.16–8.04 (m, 2H, H–3', H–6'), 7.80 (m, 1H, H–4'), 7.68 (m, 1H, CONH), 3.74 (m, 1H, H–2), 3.04 (m, 1H, N–CH–1''), 2.04 (m, 2H, H ₂ –4), 1.82 (m, 1H, H _A –3), 1.64 (m, 1H, H _B –3), 0.95–0.88 (m, 6H, CH ₃ –2'', CH ₃ –3'')	42.33 42.22	4.57 4.48	7.05 6.96
12	M + H ⁺ peak at m/z 412	3320, 3100 (N–H str of CONH), 3016 (Ar–C–H str), 2880 (ali C–H str), 1700 (C=O str), 1556, 1440 (ali C–H def), 1332 & 1164 (S=O str of SO ₂ NH), 975, 793 & 750 (Ar–C–H def)	12.66 (s, 1H, COOH), 8.40 (d, 1H, SO ₂ NH), 8.16–8.06 (m, 2H, H–3', H–6'), 7.78 (m, 1H, H–4'), 7.64(m, 1H, CONH), 3.70 (m, 1H, H–2), 3.04 (m, 2H, N–CH ₂ –1''), 2.18 (m, 2H, H ₂ –4), 1.92 (m, 1H, H _A –3), 1.72 (m, 1H, H _B –3), 1.42–1.24 (m, 4H, CH ₂ –2'', CH ₂ –3''), 0.98 (m, 3H, CH ₃ –4'')	43.80 43.70	4.90 4.84	6.81 6.75
13	M + H ⁺ peak at m/z 412	3314, 3112 (N–H str of CONH), 3020 (Ar–C–H str), 2872 (ali C–H str), 1705 (C=O str), 1558, 1438 (ali C–H def), 1334 & 1162 (S=O str of SO ₂ NH), 975, 798 & 752 (Ar–C–H def)	12.66 (s, 1H, COOH), 8.48 (d, 1H, SO ₂ NH), 8.22–8.12 (m, 2H, H–3', H–6'), 7.84 (m, 1H, H–4'), 7.70 (m, 1H, CONH), 3.84 (m, 1H, H–2), 3.06 (m, 2H, N–CH ₂ –1''), 2.12 (m, 2H, H ₂ –4), 1.86 (m, 1H, H _A –3), 1.72 (m, 1H, H _B –3), 1.42 (m, 1H, CH–2''), 1.15–0.98 (m, 6H, CH ₃ –3'', CH ₃ –4'')	43.80 43.65	4.90 4.80	6.81 6.78

Table 3. (Continued)

Cpd	Mass (FAB)	IR (KBr, cm ⁻¹)	¹ HNMR (300 MHz, DMSO- <i>d</i> ₆)	C,H,N: %calcd/found		
				C	H	N
14	M + H ⁺ peak at m/z 440	3318, 3105 (N–H str of CONH), 3012 (Ar–C–H str), 2876 (ali C–H str), 1705 (C=O str), 1556, 1444 (ali C–H def), 1334 & 1162 (S=O str of SO ₂ NH), 970, 790 & 748 (Ar–C–H def)	12.62 (s, 1H, COOH), 8.46 (d, 1H, SO ₂ NH), 8.12–8.02 (m, 2H, H–3', H–6'), 7.72 (m, 1H, H–4'), 7.60 (m, 1H, CONH), 3.68 (m, 1H, H–2), 3.00 (m, 2H, N–CH ₂ –1''), 2.12 (m, 2H, H ₂ –4), 1.90 (m, 1H, H _A –3), 1.68 (m, 1H, H _B –3), 1.40–1.16 (m, 8H, CH ₂ –2'', CH ₂ –3'', CH ₂ –4'', CH ₂ –5''), 0.90 (m, 3H, CH ₃ –6'')	46.47 46.32	5.51 5.43	6.38 6.29
15	M + H ⁺ peak at m/z 438	3308, 3102 (N–H str of CONH), 3028 (Ar–C–H str), 2878 (ali C–H str), 1698 (C=O str), 1560, 1440 (ali C–H def), 1336 & 1162 (S=O str of SO ₂ NH), 972, 794 & 750 (Ar–C–H def)	12.62 (s, 1H, COOH), 8.48 (d, 1H, SO ₂ NH), 8.22–8.12 (m, 2H, H–3', H–6'), 7.82 (m, 1H, H–4'), 7.68 (m, 1H, CONH), 3.74m, 1H, H–2), 2.12 (m, 2H, H ₂ –4), 1.80 (m, 1H, H _A –3), 1.62 (m, 1H, H _B –3) 1.38–1.10 (m, 11H, cyclohexyl protons)	46.69 46.58	5.07 5.02	6.41 6.54
16	M + H ⁺ peak at m/z 432	3322, 3116 (N–H str of CONH), 3020 (Ar–C–H str), 2878 (ali C–H str), 1702 (C=O str), 1548, 1442 (ali C–H def), 1336 & 1160 (S=O str of SO ₂ NH), 972, 796 & 748 (Ar–C–H def)	12.62 (s, 1H, COOH), 8.38 (d, 1H, SO ₂ NH), 8.10–8.02 (m, 2H, H–3', H–6'), 7.84 (m, 5H, ph–protons), 7.70 (m, 1H, H–4'), 7.58 (m, 1H, CONH), 3.74 (m, 1H, H–2), 2.12 (m, 2H, H ₂ –4), 1.86 (m, 1H, H _A –3), 1.72 (m, 1H, H _B –3)	47.34 47.13	3.74 3.68	6.50 6.56
17	M + H ⁺ peak at m/z 446	3316, 3106 (N–H str of CONH), 3012 (Ar–C–H str), 2876 (ali C–H str), 1700 (C=O str), 1556, 1440 (ali C–H def), 1332 & 1160 (S=O str of SO ₂ NH), 977, 798 & 748 (Ar–C–H def)	12.66 (s, 1H, COOH), 8.46 (d, 1H, SO ₂ NH), 8.20–8.08 (m, 2H, H–3', H–6'), 7.88–7.76 (m, 6H, H–4', ph–protons), 7.72 (m, 1H, CONH), 4.22 (m, 2H, CH ₂ –ph), 3.76 (m, 1H, H–2), 2.70 (s, 3H, Ar–CH ₃), 2.10 (m, 2H, H ₂ –4), 1.92 (m, 1H, H _A –3), 1.66 (m, 1H, H _B –3)	48.55 48.69	4.07 4.17	6.29 6.32
18	M + H ⁺ peak at m/z 484	3028 (Ar–C–H str), 2836 (ali C–H str), 1706 (C=O str), 1565, 1448 (ali C–H def), 1326 & 1160 (S=O str of SO ₂ NH), 978, 796 & 754 (Ar–C–H def)	12.64 (s, 1H, COOH), 8.52 (d, 1H, SO ₂ NH), 8.22–7.96 (m, 3H, H–3', H–4', H–6'), 3.74 (m, 1H, H–2), 3.18–3.00 (m, 6H, CH ₃ –1'', CH ₃ –2''), 2.08 (m, 2H, H ₂ –4), 1.88 (m, 1H, H _A –3), 1.62 (m, 1H, H _B –3)	40.74 40.58	4.25 4.21	7.31 7.42
19	M + H ⁺ peak at m/z 440	3024 (Ar–C–H str), 2830 (ali C–H str), 1700 (C=O str), 1562, 1440 (ali C–H def), 1328 & 1164 (S=O str of SO ₂ NH), 976, 794 & 748 (Ar–C–H def)	12.70 (s, 1H, COOH), 8.56 (d, 1H, SO ₂ NH), 8.26–7.16 (m, 3H, H–3', H–4', H–6'), 3.68 (m, 1H, H–2), 3.16 (m, 2H, N–CH–1'', N–CH–2''), 2.18 (m, 2H, H ₂ –4), 1.96 (m, 1H, H _A –3), 1.72 (m, 1H, H _B –3), 1.54–1.32 (m, 12H, CH ₃ –3'', CH ₃ –4'', CH ₃ –5'', CH ₃ –6'')	46.47 46.45	5.51 5.48	6.38 6.32
20	M + H ⁺ peak at m/z 315	3320, 3106 (N–H str of CONH), 3022 (Ar–C–H str), 2878 (ali C–H str), 1702 (C=O str), 1552, 1440 (ali C–H def), 1336 & 1164 (S=O str of SO ₂ NH), 972, 794 & 750 (Ar–C–H def)	12.70 (s, 1H, COOH), 8.46 (d, 1H, SO ₂ NH), 8.20–8.12 (m, 3H, H–3', H–5', H–6'), 7.70 (m, 2H, CONH ₂), 3.76 (m, 1H, H–2), 2.70–2.62 (m, 6H, Ar–CH ₃ –2', Ar–CH ₃ –4'), 2.12 (m, 2H, H ₂ –4), 1.88 (m, 1H, H _A –3), 1.72 (m, 1H, H _B –3)	49.56 49.67	5.77 5.72	8.91 8.83
21	M + H ⁺ peak at m/z 371	3322, 3110 (N–H str of CONH), 3018 (Ar–C–H str), 2874 (ali C–H str), 1705 (C=O str), 1552, 1444 (ali C–H def), 1332 & 1160 (S=O str of SO ₂ NH), 974, 796 & 752 (Ar–C–H def)	12.66 (s, 1H, COOH), 8.52 (d, 1H, SO ₂ NH), 8.22–8.16 (m, 3H, H–3', H–5', H–6'), 7.64 (m, 1H, CONH), 3.72 (m, 1H, H–2), 2.70–2.60 (m, 6H, Ar–CH ₃ –2', Ar–CH ₃ –4'), 2.14 (m, 2H, H ₂ –4), 1.86 (m, 1H, H _A –3), 1.70 (m, 1H, H _B –3)	55.12 55.02	7.07 6.98	7.56 7.62
22	M + H ⁺ peak at m/z 371	3016 (Ar–C–H str), 2872 (ali C–H str), 1700 (C=O str), 1554, 1446 (ali C–H def), 1330 & 1162 (S=O str of SO ₂ NH), 976, 798 & 750 (Ar–C–H def)	12.72 (s, 1H, COOH), 8.48 (d, 1H, SO ₂ NH), 8.18–8.10 (m, 3H, H–3', H–5', H–6'), 3.70 (m, 1H, H–2), 3.16–3.04 (m, 4H, N–CH ₂ –1'', N–CH ₂ –2''), 2.68–2.56 (m, 6H, Ar–CH ₃ –2', Ar–CH ₃ –4'), 2.12 (m, 2H, H ₂ –4), 1.80 (m, 1H, H _A –3), 1.73 (m, 1H, H _B –3), 1.60–1.46 (m, 6H, CH ₃ –3'', CH ₃ –4'')	55.12 54.98	7.07 7.02	7.56 7.48
23	M + H ⁺ peak at m/z 343	3322, 3118 (N–H str of CONH), 3026 (Ar–C–H str), 2868 (ali C–H str), 1705 (C=O str), 1564, 1444 (ali C–H def), 1332 & 1160 (S=O str of SO ₂ NH), 975, 794 & 750 (Ar–C–H def)	12.64 (s, 1H, COOH), 8.46 (d, 1H, SO ₂ NH), 8.28–7.98 (m, 4H, H–2', H–3', H–5', H–6'), 7.68 (m, 2H, CONH ₂), 3.66 (m, 1H, H–2), 2.16 (m, 2H, H ₂ –4), 2.08–1.92 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 1.82 (m, 1H, H _A –3), 1.70 (m, 1H, H _B –3)	52.62 52.57	6.48 6.37	8.18 8.10

Table 3. (Continued)

Cpd	Mass (FAB)	IR (KBr, cm ⁻¹)	¹ HNMR (300 MHz, DMSO- <i>d</i> ₆)	C,H,N: %calcd/found		
				C	H	N
24	M + H ⁺ peak at m/z 357	3316, 3112 (N–H str of CONH), 3022 (Ar–C–H str), 2856 (ali C–H str), 1695 (C=O str), 1558, 1440 (ali C–H def), 1330 & 1163 (S=O str of SO ₂ NH), 973, 797 & 754 (Ar–C–H def)	12.68 (s, 1H, COOH), 8.50 (d, 1H, SO ₂ NH), 8.26–8.05 (m, 4H, H–2', H–3', H–5', H–6'), 7.62 (m, 1H, CONH), 3.69 (m, 1H, H–2), 3.12 (m, 3H, CH ₃ –1''), 2.16 (m, 2H, H ₂ –4), 2.08–1.94 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 1.82 (m, 1H, H _A –3), 1.68 (m, 1H, H _B –3)	53.91 53.93	6.79 6.78	7.86 7.82
25	M + H ⁺ peak at m/z 371	3320, 3115 (N–H str of CONH), 3025 (Ar–C–H str), 2856 (ali C–H str), 1698(C=O str), 1560, 1445 (ali C–H def), 1335 & 1168 (S=O str of SO ₂ NH), 975, 801 & 757 (Ar–C–H def)	12.71 (s, 1H, COOH), 8.55 (d, 1H, SO ₂ NH), 8.24–8.10 (m, 4H, H–2', H–3', H–5', H–6'), 7.67(m, 1H, CONH), 3.73 (m, 1H, H–2), 3.16 (m, 2H, N–CH ₂ –1''), 2.19 (m, 2H, H ₂ –4), 2.10–1.96 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 1.85 (m, 1H, H _A –3), 1.72 (m, 1H, H _B –3), 1.64 (m, 3H, CH ₃ –2'')	55.12 55.20	7.07 7.16	7.56 7.68
26	M + H ⁺ peak at m/z 385	3327, 3119 (N–H str of CONH), 3029 (Ar–C–H str), 2858 (ali C–H str), 1700 (C=O str), 1566, 1449 (ali C–H def), 1339 & 1170 (S=O str of SO ₂ NH), 978, 796 & 758 (Ar–C–H def)	12.75 (s, 1H, COOH), 8.60 (d, 1H, SO ₂ NH), 8.27–8.15 (m, 4H, H–2', H–3', H–5', H–6'), 7.70(m, 1H, CONH), 3.78 (m, 1H, H–2), 3.20 (m, 2H, N–CH ₂ –1''), 2.26 (m, 2H, H ₂ –4), 2.14–1.98 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 1.89(m, 1H, H _A –3), 1.75 (m, 1H, H _B –3), 1.66 (m, 3H, CH ₃ –2'')	56.23 56.65	7.34 7.23	7.29 7.33
27	M + H ⁺ peak at m/z 385	3319, 3112 (N–H str of CONH), 3028 (Ar–C–H str), 2855 (ali C–H str), 1695(C=O str), 1562, 1448 (ali C–H def), 1340 & 1170(S=O str of SO ₂ NH), 975, 799 & 754 (Ar–C–H def)	12.68 (s, 1H, COOH), 8.58 (d, 1H, SO ₂ NH), 8.23–8.11 (m, 4H, H–2', H–3', H–5', H–6'), 7.68(m, 1H, CONH), 3.75 (m, 1H, H–2), 3.18 (m, 1H, N–CH–1''), 2.20 (m, 2H, H ₂ –4), 2.10–1.96 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 1.88 (m, 1H, H _A –3), 1.77 (m, 1H, H _B –3), 1.63–1.50 (m, 6H, CH ₃ –2'', CH ₃ –3'')	56.23 56.27	7.34 7.38	7.29 7.12
28	M + H ⁺ peak at m/z 399	3324, 3109 (N–H str of CONH), 3029 (Ar–C–H str), 2853 (ali C–H str), 1692 (C=O str), 1564, 1451 (ali C–H def), 1343 & 1169 (S=O str of SO ₂ NH), 972, 802 & 749 (Ar–C–H def)	12.65 (s, 1H, COOH), 8.52 (d, 1H, SO ₂ NH), 8.20–8.13 (m, 4H, H–2', H–3', H–5', H–6'), 7.72 (m, 1H, CONH), 3.77(m, 1H, H–2), 3.15 (m, 2H, N–CH ₂ –1''), 2.32 (m, 2H, H ₂ –4), 2.16–2.02 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 1.92 (m, 1H, H _A –3), 1.82 (m, 1H, H _B –3), 1.72–1.52 (m, 7H, CH ₂ –2'', CH ₂ –3'', CH ₃ –4'')	57.26 57.12	7.59 7.53	7.03 6.95
29	M + H ⁺ peak at m/z 399	3324, 3112 (N–H str of CONH), 3025 (Ar–C–H str), 2850 (ali C–H str), 1698 (C=O str), 1568, 1448 (ali C–H def), 1340 & 1164 (S=O str of SO ₂ NH), 971, 798 & 752 (Ar–C–H def)	12.72 (s, 1H, COOH), 8.56 (d, 1H, SO ₂ NH), 8.18–8.09 (m, 4H, H–2', H–3', H–5', H–6'), 7.78 (m, 1H, CONH), 3.80 (m, 1H, H–2), 3.17 (m, 2H, N–CH ₂ –1''), 2.35 (m, 2H, H ₂ –4), 2.16–1.98 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 1.89 (m, 1H, H _A –3), 1.78 (m, 1H, H _B –3), 1.68–1.52 (m, 7H, CH–2'', CH ₃ –3'', CH ₃ –4'')	57.26 57.06	7.59 7.50	7.03 6.98
30	M + H ⁺ peak at m/z 427	3320, 3114 (N–H str of CONH), 3022 (Ar–C–H str), 2856 (ali C–H str), 1705 (C=O str), 1556, 1442 (ali C–H def), 1332 & 1160 (S=O str of SO ₂ NH), 973, 794 & 748 (Ar–C–H def)	12.66 (s, 1H, COOH), 8.52 (d, 1H, SO ₂ NH), 8.14–8.06 (m, 4H, H–2', H–3', H–5', H–6'), 7.81 (m, 1H, CONH), 3.78 (m, 1H, H–2), 3.12 (m, 2H, N–CH ₂ –1''), 2.38 (m, 2H, H ₂ –4), 2.18–2.06 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 1.92 (m, 1H, H _A –3), 1.82 (m, 1H, H _B –3), 1.74–1.48 (m, 11H, CH ₂ –2'', CH ₂ –3'', CH ₂ –4'', CH ₂ –5'', CH ₃ –6'')	59.13 58.96	8.03 7.92	6.57 6.48
31	M + H ⁺ peak at m/z 425	3321, 3115 (N–H str of CONH), 3027 (Ar–C–H str), 2855 (ali C–H str), 1701 (C=O str), 1558, 1440 (ali C–H def), 1336 & 1164 (S=O str of SO ₂ NH), 976, 797 & 752 (Ar–C–H def)	12.72 (s, 1H, COOH), 8.56 (d, 1H, SO ₂ NH), 8.18–8.06 (m, 4H, H–2', H–3', H–5', H–6'), 7.85 (m, 1H, CONH), 3.82 (m, 1H, H–2), 2.44 (m, 2H, H ₂ –4), 2.22–2.12 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 1.96 (m, 1H, H _A –3), 1.86 (m, 1H, H _B –3), 1.52–1.26 (m, 11H, cyclohexyl protons)	59.41 59.28	7.60 7.54	6.60 6.62

Table 3. (Continued)

Cpd	Mass (FAB)	IR (KBr, cm ⁻¹)	¹ HNMR (300 MHz, DMSO- <i>d</i> ₆)	C,H,N: %calcd/found		
				C	H	N
32	M + H ⁺ peak at m/z 419	3319, 3114 (N-H str of CONH), 3023 (Ar-C-H str), 2857 (ali C-H str), 1699 (C=O str), 1559, 1445 (ali C-H def), 1337 & 1165 (S=O str of SO ₂ NH), 977, 798 & 753 (Ar-C-H def)	12.64 (s, 1H, COOH), 8.46 (d, 1H, SO ₂ NH), 8.12–8.02 (m, 4H, H-2', H-3', H-5', H-6'), 7.88 (m, 1H, CONH), 7.72 (m, 5H, ph.-protons), 3.86 (m, 1H, H- 2), 2.44 (m, 2H, H ₂ -4), 2.22–2.08 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 1.94 (m, 1H, H _A -3), 1.88 (m, 1H, H _B -3)	60.27 60.08	6.26 6.14	6.69 6.54
33	M + H ⁺ peak at m/z 433	3324, 3118 (N-H str of CONH), 3028 (Ar-C-H str), 2855 (ali C-H str), 1703 (C=O str), 1556, 1442 (ali C-H def), 1332 & 1162 (S=O str of SO ₂ NH), 975, 794 & 750 (Ar-C-H def)	12.68 (s, 1H, COOH), 8.52 (d, 1H, SO ₂ NH), 8.18–8.08 (m, 4H, H-2', H-3', H-5', H-6'), 7.92 (m, 1H, CONH), 7.74–7.66 (m, 5H, ph.-protons), 4.28 (m, 2H, CH ₂ -ph), 3.84 (m, 1H, H-2), 2.46 (m, 2H, H ₂ -4), 2.20–2.12 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 2.02 (m, 1H, H _A -3), 1.94 (m, 1H, H _B -3)	61.09 60.92	6.52 6.42	6.48 6.36
34	M + H ⁺ peak at m/z 371	3025 (Ar-C-H str), 2858 (ali C-H str), 1706 (C=O str), 1548, 1440 (ali C-H def), 1336 & 1164 (S=O str of SO ₂ NH), 977, 796 & 754 (Ar-C-H def)	12.62 (s, 1H, COOH), 8.55 (d, 1H, SO ₂ NH), 8.14–8.06 (m, 4H, H-2', H-3', H-5', H-6'), 3.80 (m, 1H, H-2), 3.24–3.12 (m, 6H, CH ₃ -1'', CH ₃ -2''), 2.40 (m, 2H, H ₂ -4), 2.10–1.98 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 1.92 (m, 1H, H _A -3), 1.82 (m, 1H, H _B -3)	55.12 54.98	7.07 6.99	7.56 7.48
35	M + H ⁺ peak at m/z 399	3025 (Ar-C-H str), 2858 (ali C-H str), 1706 (C=O str), 1548, 1440 (ali C-H def), 1336 & 1164 (S=O str of SO ₂ NH), 977, 796 & 754 (Ar-C-H def)	12.68 (s, 1H, COOH), 8.50 (d, 1H, SO ₂ NH), 8.10–8.02 (m, 4H, H-2', H-3', H-5', H-6'), 3.74 (m, 1H, H-2), 3.22–3.08 (m, 4H, CH ₂ -1'', CH ₂ -2''), 2.44 (m, 2H, H ₂ -4), 2.18–2.04 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 1.98 (m, 1H, H _A -3), 1.80 (m, 1H, H _B -3), 1.72–1.56 (m, 6H, CH ₃ -3'', CH ₃ -4'')	57.26 57.22	7.59 7.49	7.03 6.98
36	M + H ⁺ peak at m/z 427	3020 (Ar-C-H str), 2852 (ali C-H str), 1700 (C=O str), 1552, 1444 (ali C-H def), 1332 & 1160 (S=O str of SO ₂ NH), 975, 794 & 752 (Ar-C-H def)	12.62 (s, 1H, COOH), 8.56 (d, 1H, SO ₂ NH), 8.14–8.08 (m, 4H, H-2', H-3', H-5', H-6'), 3.78 (m, 1H, H-2), 3.20–3.12 (m, 2H, N-CH-1'', N-CH-2''), 2.42 (m, 2H, H ₂ -4), 2.22–2.08 (m, 9H, three CH ₃ of <i>t</i> -Butyl), 2.02 (m, 1H, H _A -3), 1.94 (m, 1H, H _B -3), 1.74–1.62 (m, 12H, CH ₃ -3'', CH ₃ -4'', CH ₃ -5'', CH ₃ -6'')	59.13 59.02	8.03 7.92	6.57 6.62

2.2 Biological Activity

The title compounds were evaluated for their possible anticancer activity *in vivo* against Ehrlich Ascites Carcinoma (EAC) cell in Swiss Albino mice as per the standard procedure [13–15, 17] using percentage of tumor weight inhibition (% TWI) as the biological activity data.

2.2.1 Tumor cells

EAC were maintained *in vivo* in Swiss Albino mice, by passaging every 10 days. EAC cells of 9 day old were used for the screening of the entire title compounds.

2.2.2 Animals

Swiss albino mice (either sex) of 10 weeks old with an average body weight of 18–20 grams were used. All mice were kept on basal metabolic diet with water *ad libitum*.

2.2.3 Procedure

Two groups of Swiss Albino Mice, each containing 5 healthy animal of the same sex, approximately of the same age and body weight, were selected at random and kept in two different cages under identical conditions. One of these two groups served as control while the other as test.

Ehrlich Ascites Carcinoma (EAC) cells were collected from the donor mice and were suspended in sterile isotonic solution (0.9% w/v NaCl). A definite number (about 2×10^6 cells/0.2 ml) of these living viable cells was injected or implanted into the peritoneal cavity of each mouse. A day of incubation was allowed to establish the disease in the body before the start of the drug administration. From the second day of transplantation up to the eighth day, a suitable challenge dose (0.2 mM/kg body weight) of the drug solution/suspension in sterile phosphate buffer (pH 7.2) was injected intraperitoneally to each mouse in the test group at 24 hr interval. Thus, seven doses of the drug were administered to each mouse in the test group. On the ninth day, food and water was withheld 18 hr before the start of the testing operation. The weights of all the animals were recorded before they were sacrificed. The peritoneal cavity was dissected and by a syringe, the ascitic fluid was withdrawn to a suitable volume, collected in sterile ice-cold saline and preserved in ice bath. The fluid was sucked by adsorbent cotton. The weight of the 5 mice after sacrifice was recorded. The evaluation of the test drug was made by comparing the tumor weight of the test with that of the control.

2.3 QSAR Methodology

2.3.1 Dataset and parameter

Antitumor activities of thirty-two 5-*N*-substituted-2-(substituted benzenesulphonyl) glutamines of the general structure shown in Figure 1 were used to develop QSAR models separately. Percentage tumor weight inhibition (%TWI) has been considered as biological activity parameters for QSAR studies. All these activities are calibrated to the logarithmic value.

2.3.2 Physicochemical parameters

The physicochemical parameters like, hydrophobic constant π , electronic parameters σ (Hammett constant), \mathcal{R} (resonance effect) & \mathcal{F} (field effect), MR (Molar Refractivity), steric parameter E_s , sterimol parameters like L , B_1 , B_5 were collected from the literature [21, 22] and are listed in Table 5. σ , \mathcal{R} , \mathcal{F} all these parameters describe the influence of a certain group or substituent on electron density distribution. MR is largely a measure of volume with a small correction for polarizability. MR values were scaled by a factor 0.1. E_s is the classical Taft parameter derived from the rate of hydrolysis of aliphatic esters.

2.3.3 Electrotopological state atom (ETSA) indices

The ETA index of each atom explained electronic and topological information of all other atoms within the structure [23, 24]. ETA indices were calculated using the computer program 'Mouse' [25] developed in our laboratory. For calculation of E-state index, arbitrary numbering was used and these are shown in Figure 2.




Figure 2. General Structure of 5-*N*-substituted-2-(substituted benzenesulphonyl) glutamines (5–36) with arbitrary numbers used for ETSA calculations.

2.3.4 Statistical Analysis

2.3.4.1 Correlation Analysis

Correlation analysis [26] of biological activity, ETSA indices and physicochemical parameters was carried out. Inter-correlated parameters were eliminated stepwise depending on their individual correlation with the biological activity. All possible combinations of parameters were considered for multiple regression analysis.

2.3.4.2 Multiple Regression Analysis

Multiple regression analysis [26–29] was carried out by ‘Multi Regress’ [30], a program developed in our laboratory. The statistical quality of the regression equation were justified by parameters like squared correlation coefficient (R^2), adjusted R^2 (R^2_A), variance ratio (F), standard error of estimate (s). All the final equations have significant regression coefficients, intercepts and variance ratio (F) and that are more than 95% level. Use of more than one variable in the multivariate equation was justified by autocorrelation study.

2.3.4.3 Cross Validation

The predictive powers of the equation were validated by Leave-One-Out (LOO) cross-validation method [31]. Predicted residual sum of square ($PRESS$), total sum of squares (SSY), cross-validated R^2 (R^2_{CV}), standard error of $PRESS$ (S_{PRESS}) and standard deviation of the error of prediction (S_{DEP}) for the QSAR equations were considered for the validation of the models.

3 RESULTS AND DISCUSSION

3.1 Synthesis

32 new 5-*N*-substituted-2-(substituted benzenesulphonyl) glutamines were synthesized. There were four types of substitutions at aromatic ring whereas the aliphatic side chain contains thirteen substitutions. Percentage of yields of the final compounds was ranging from 45 to 92% as shown in Table 2.

3.2 Screening

All the final compounds were screened for their possible anticancer activity against Ehrlich Ascites Carcinoma (EAC) cell as per standard method [13–15, 17]. Compounds were dissolved or suspended in PBS with 2% Tween 80 (whenever necessary) separately. The solution or suspension of the test compounds was administered at a dose level of 2 mmol/kg/day intraperitoneally (i.p.) for 7 consecutive days. % Inhibition of tumor weight was considered as the biological activity parameter for QSAR study and thus, all the activity data are converted into logarithmic scale. Table 4 shows the anticancer activity of all the compounds.

Table 4. Anticancer activities of 5-*N*-Substituted-2-(substituted benzenesulphonyl) glutamines 5–36

Cpd	R ₁	R ₂	R ₃	R ₄	R ₅	%TWI	Log (TWI)
5	H	Br	H	<i>i</i> -C ₃ H ₇	H	46.51	1.668
6	H	Br	H	<i>c</i> -C ₆ H ₁₁	H	63.24	1.801
7	H	Br	H	C ₆ H ₅ CH ₂	H	34.39	1.536
8	H	Br	H	CH ₃	CH ₃	26.56	1.424
9	Cl	H	Cl	CH ₃	H	40.28	1.605
10	Cl	H	Cl	<i>n</i> -C ₃ H ₇	H	40.00	1.602
11	Cl	H	Cl	<i>i</i> -C ₃ H ₇	H	52.15	1.717
12	Cl	H	Cl	<i>n</i> -C ₄ H ₉	H	47.67	1.678
13	Cl	H	Cl	<i>i</i> -C ₄ H ₉	H	48.21	1.683
14	Cl	H	Cl	<i>n</i> -C ₆ H ₁₃	H	56.36	1.751
15	Cl	H	Cl	<i>c</i> -C ₆ H ₁₁	H	56.09	1.749
16	Cl	H	Cl	C ₆ H ₅	H	48.19	1.683
17	Cl	H	Cl	C ₆ H ₅ CH ₂	H	42.76	1.631
18	Cl	H	Cl	CH ₃	CH ₃	22.28	1.348
19	Cl	H	Cl	<i>i</i> -C ₃ H ₇	<i>i</i> -C ₃ H ₇	44.23	1.646
20	CH ₃	CH ₃	H	H	H	23.23	1.366
21	CH ₃	CH ₃	H	<i>n</i> -C ₄ H ₉	H	35.55	1.551
22	CH ₃	CH ₃	H	C ₂ H ₅	C ₂ H ₅	21.86	1.340
23	H	<i>t</i> -C ₄ H ₉	H	H	H	55.01	1.300
24	H	<i>t</i> -C ₄ H ₉	H	CH ₃	H	40.01	1.602
25	H	<i>t</i> -C ₄ H ₉	H	C ₂ H ₅	H	32.38	1.510
26	H	<i>t</i> -C ₄ H ₉	H	<i>n</i> -C ₃ H ₇	H	32.81	1.516
27	H	<i>t</i> -C ₄ H ₉	H	<i>i</i> -C ₃ H ₇	H	45.68	1.660
28	H	<i>t</i> -C ₄ H ₉	H	<i>n</i> -C ₄ H ₉	H	45.16	1.655
29	H	<i>t</i> -C ₄ H ₉	H	<i>i</i> -C ₄ H ₉	H	38.55	1.586
30	H	<i>t</i> -C ₄ H ₉	H	<i>n</i> -C ₆ H ₁₃	H	50.82	1.706
31	H	<i>t</i> -C ₄ H ₉	H	<i>c</i> -C ₆ H ₁₁	H	61.42	1.788
32	H	<i>t</i> -C ₄ H ₉	H	C ₆ H ₅	H	67.85	1.832
33	H	<i>t</i> -C ₄ H ₉	H	C ₆ H ₅ CH ₂	H	33.93	1.531
34	H	<i>t</i> -C ₄ H ₉	H	CH ₃	CH ₃	23.57	1.372
35	H	<i>t</i> -C ₄ H ₉	H	C ₂ H ₅	C ₂ H ₅	22.79	1.358
36	H	<i>t</i> -C ₄ H ₉	H	<i>i</i> -C ₃ H ₇	<i>i</i> -C ₃ H ₇	28.74	1.458

3.3 QSAR Study

To identify the chemical structural features required for antitumor activity of 5-*N*-substituted-2-(substituted benzenesulphonyl) glutamines, Quantitative Structure–Activity Relationship (QSAR) studies were performed as a part of our composite program of rational drug design [12–17,31–39]. The physicochemical parameters and ETSA indices were used to develop QSAR models. Important

physicochemical parameters and ETSA indices are shown in Table 5. Correlation analysis was performed using Log (TWI) as the dependent variable and physicochemical and ETSA indices as independent parameter. The result of this analysis is shown in Table 6.

Table 5. Physicochemical parameters and ETSA indices of glutamine analogs 5–36

Cpd	πR_1	πR_2	πR_4	MRR_4	LR_4	B_1R_4	S_2	S_3	S_4	S_5	S_6	S_7	S_{AV}
5	0.000	-0.170	0.870	1.287	4.110	1.900	1.542	0.706	1.542	1.358	-0.042	-3.975	0.226
6	0.000	-0.170	2.070	2.497	6.170	1.910	1.557	0.714	1.557	1.376	-0.028	-3.979	0.239
7	0.000	-0.170	2.060	2.867	4.620	1.520	1.543	0.701	1.543	1.359	-0.050	-4.007	0.218
8	0.000	-0.170	0.040	0.357	2.870	1.520	1.541	0.707	1.541	1.357	-0.040	-3.965	0.228
9	0.410	0.000	0.040	0.357	2.870	1.520	1.290	1.406	0.142	1.115	-0.325	-4.190	-0.112
10	0.410	0.000	1.090	1.290	4.920	1.520	1.298	1.412	0.145	1.124	-0.320	-4.201	-0.108
11	0.410	0.000	0.870	1.287	4.110	1.900	1.296	1.41	0.142	1.121	-0.340	-4.207	-0.116
12	0.410	0.000	1.620	1.747	6.170	1.520	1.301	1.415	0.146	1.127	-0.318	-4.205	-0.107
13	0.410	0.000	1.490	1.747	4.920	1.520	1.299	1.413	0.144	1.125	-0.321	-4.209	-0.110
14	0.410	0.000	2.680	2.677	8.220	1.520	1.306	1.419	0.148	1.133	-0.316	-4.211	-0.104
15	0.410	0.000	2.070	2.497	6.170	1.910	1.311	1.424	0.152	1.140	-0.310	-4.211	-0.099
16	0.410	0.000	2.270	2.277	6.280	1.520	1.293	1.408	0.134	1.118	-0.336	-4.243	-0.125
17	0.410	0.000	2.060	2.867	4.620	1.520	1.297	1.411	0.137	1.122	-0.331	-4.239	-0.121
18	0.410	0.000	0.040	0.357	2.870	1.520	1.295	1.41	0.144	1.121	-0.322	-4.197	-0.110
19	0.410	0.000	0.870	1.287	4.110	1.900	1.306	1.419	0.146	1.134	-0.318	-4.239	-0.110
20	-0.040	-0.130	0.000	0.103	2.060	1.000	1.688	0.899	1.626	1.416	0.013	-3.982	0.332
21	-0.040	-0.130	1.620	1.747	6.170	1.520	1.714	0.913	1.651	1.448	0.039	-3.982	0.357
22	-0.040	-0.130	0.560	0.817	4.110	1.520	1.716	0.96	1.654	1.451	0.034	-3.999	0.363
23	0.000	-0.130	0.000	0.103	2.060	1.000	1.688	0.959	1.688	1.426	-0.031	-3.997	0.347
24	0.000	-0.130	0.040	0.357	2.870	1.520	1.703	0.969	1.703	1.445	-0.011	-3.983	0.365
25	0.000	-0.130	0.560	0.817	4.110	1.520	1.707	0.971	1.707	1.450	-0.009	-3.989	0.367
26	0.000	-0.130	1.090	1.290	4.920	1.520	1.711	0.972	1.712	1.454	-0.006	-3.993	0.370
27	0.000	-0.130	0.870	1.287	4.110	1.900	1.709	0.97	1.709	1.452	-0.010	-3.999	0.366
28	0.000	-0.130	1.620	1.747	6.170	1.520	1.714	0.973	1.714	1.457	-0.005	-3.997	0.371
29	0.000	-0.130	1.490	1.747	4.920	1.520	1.712	0.971	1.712	1.455	-0.007	-4.002	0.368
30	0.000	-0.130	2.680	2.677	8.220	1.520	1.718	0.975	1.718	1.463	-0.002	-4.004	0.374
31	0.000	-0.130	2.070	2.497	6.170	1.910	1.724	0.978	1.724	1.470	-0.004	-4.003	0.378
32	0.000	-0.130	2.270	2.277	6.280	1.710	1.706	0.963	1.706	1.448	-0.022	-4.035	0.353
33	0.000	-0.130	2.060	2.867	4.620	1.520	1.710	0.965	1.71	1.453	-0.017	-4.032	0.358
34	0.000	-0.130	0.040	0.357	2.870	1.520	1.708	0.971	1.708	1.451	-0.008	-3.990	0.368
35	0.000	-0.130	0.560	0.817	4.110	1.520	1.716	0.974	1.716	1.461	-0.002	-4.001	0.373
36	0.000	-0.130	0.870	1.287	4.110	1.900	1.719	0.972	1.719	1.464	-0.005	-4.022	0.369

Multiple regression analysis using the combination of physicochemical parameters πR_4 (hydrophobicity of the R_4 substituent), B_1R_4 (sterimol parameter for minimum width of R_4 substituent) developed the following QSAR model as shown bellow:

$$\begin{aligned} \text{Log (TWI)} &= 1.024 (\pm 0.111) + 0.103 (\pm 0.019) \pi R_4 + 0.274 (\pm 0.072) B_1R_4 \\ n &= 32 \quad R^2 = 0.679 \quad R^2_A = 0.657 \quad F_{(2,29)} = 30.664 \quad p < 0.001 \quad s = 0.087 \\ SSY &= 0.688 \quad PRESS = 0.264 \quad R^2_{CV} = 0.616 \quad S_{PRESS} = 0.095 \quad S_{DEP} = 0.091 \end{aligned} \quad (1)$$

where n is the number of data points, R^2 , R^2_A , F , p , s are the squared correlation coefficient, adjusted R^2 , ratio between the variances of observed and calculated activities, probability factor related to the F -ratio, standard error of estimate respectively. SSY , $PRESS$, R^2_{CV} , S_{PRESS} , S_{DEP} are total sum of squares, predicted residual sum of square, cross-validated R^2 , standard error of $PRESS$ and standard deviation of the error of prediction respectively. This model explains 67.90 % of the variances in the activity data. Eq. (1) suggests the importance of B_1R_4 and πR_4 in tumor weight inhibition of the glutamine analogs. Positive coefficients of B_1R_4 and πR_4 indicate that these are conducive to the

activity.

Table 6. Correlation matrix for the anticancer activity, physicochemical parameters, ETSA and indicator parameter

	\mathcal{R}_1	\mathcal{R}_2	πR_4	MRR_4	LR_4	B_1R_4	S_2	S_3	S_4	S_5
\mathcal{R}_1	1.00	0.98	0.15	0.15	0.15	0.13	-0.96	0.94	-0.99	-0.98
\mathcal{R}_2		1.00	0.13	0.11	0.15	0.06	-0.89	0.99	-0.97	-0.93
πR_4			1.00	0.97	0.91	0.30	-0.13	0.13	-0.14	-0.12
MRR_4				1.00	0.81	0.37	-0.15	0.10	-0.14	-0.12
LR_4					1.00	0.33	-0.10	0.16	-0.13	-0.09
B_1R_4						1.00	-0.15	0.06	-0.12	-0.11
S_2							1.00	-0.82	0.98	0.99
S_3								1.00	-0.92	-0.86
S_4									1.00	0.99
S_5										1.00
	S_6	S_7	S_{AV}	I	Log (TWI)					
\mathcal{R}_1	-0.99	-0.99	-0.98	-0.07	0.32					
\mathcal{R}_2	-0.99	-0.98	-0.92	-0.07	0.27					
πR_4	-0.15	-0.22	-0.14	-0.48	0.72					
MRR_4	-0.15	-0.22	-0.15	-0.45	0.69					
LR_4	-0.12	-0.19	-0.12	-0.39	0.72					
B_1R_4	-0.13	-0.14	-0.14	0.09	0.60					
S_2	0.98	0.93	0.99	0.07	-0.32					
S_3	-0.91	-0.95	-0.86	-0.06	0.26					
S_4	0.99	0.98	0.99	0.06	-0.30					
S_5	0.99	0.95	0.99	0.07	-0.30					
S_6	1.00	0.98	0.99	0.08	-0.32					
S_7		1.00	0.95	0.06	-0.33					
S_{AV}			1.00	0.07	-0.32					
I				1.00	-0.58					
Log (TWI)					1.00					

Analysis of the correlation matrix reveals that ETSA indices like S_2 , S_3 , S_4 , S_5 , S_6 , and S_7 have almost equal important contribution to the activity. But these indices are highly inter-correlated with each other and cannot be used in a single equation. Hence an average of these (S_{AV}) is considered as single best variable and incorporation of this index along with physicochemical parameters of Eq. (1) yielded Eq. (2) as shown bellow:

$$\begin{aligned} \text{Log (TWI)} &= 1.070 (\pm 0.111) + 0.099 (\pm 0.018) \pi R_4 + 0.262 (\pm 0.071) B_1R_4 - 0.121 (\pm 0.070) S_{AV} \\ n &= 32 \quad R^2 = 0.711 \quad R^2_A = 0.679 \quad F_{(3,28)} = 22.852 \quad p < 0.001 \quad s = 0.084 \\ SSY &= 0.688 \quad PRESS = 0.258 \quad R^2_{CV} = 0.625 \quad S_{PRESS} = 0.096 \quad S_{DEP} = 0.090 \end{aligned} \quad (2)$$

The percentage of explained variance of the Eq. (2) is 71.10%, whereas percentage of explained variance computed by performing LOO cross-validation is 62.50%. Negative coefficient of the S_{AV} indicates that the lower value of this index corresponds to higher anticancer activity. Inclusion of indicator parameter I for the presence of disubstitution at aliphatic side chain improve the statistical quality of the relationship as:

$$\begin{aligned} \text{Log (TWI)} &= 1.030 (\pm 0.077) + 0.054 (\pm 0.015) \pi R_4 + 0.343 (\pm 0.051) B_1 R_4 \\ &\quad - 0.113 (\pm 0.048) S_{AV} - 0.167 (\pm 0.030) I \\ n = 32 \quad R^2 &= 0.865 \quad R^2_A = 0.846 \quad F_{(4,27)} = 43.512 \quad p < 0.001 \quad s = 0.059 \\ SSY &= 0.688 \quad PRESS = 0.127 \quad R^2_{CV} = 0.815 \quad S_{PRESS} = 0.069 \quad S_{DEP} = 0.063 \end{aligned} \quad (3)$$

Eq. (3) explains 86.50% of the variances in the activity data. Negative coefficient of the 'I' also indicates that presence of disubstitution at R₄/R₅ position is disadvantageous to the activity

Another models with near statistical quality of Eq. (3) were developed using *MRR*₄ (molar refractivity of the R₄ substituent) and *LR*₄ (sterimol parameter for length of the R₄ substituent) instead of πR_4 respectively as shown in Eqs. (4) and (5):

$$\begin{aligned} \text{Log (TWI)} &= 1.029 (\pm 0.086) + 0.038 (\pm 0.016) MRR_4 + 0.351 (\pm 0.058) B_1 R_4 \\ &\quad - 0.116 (\pm 0.053) S_{AV} - 0.184 (\pm 0.033) I \\ n = 32 \quad R^2 &= 0.836 \quad R^2_A = 0.812 \quad F_{(4,27)} = 34.443 \quad p < .001 \quad s = 0.065 \\ SSY &= 0.688 \quad PRESS = 0.156 \quad R^2_{CV} = 0.773 \quad S_{PRESS} = 0.076 \quad S_{DEP} = 0.070 \end{aligned} \quad (4)$$

$$\begin{aligned} \text{Log (TWI)} &= 0.962 (\pm 0.071) + 0.032 (\pm 0.007) LR_4 + 0.333 (\pm 0.047) B_1 R_4 \\ &\quad - 0.117 (\pm 0.045) S_{AV} - 0.173 (\pm 0.026) I \\ n = 32 \quad R^2 &= 0.885 \quad R^2_A = 0.868 \quad F_{(4,27)} = 52.041 \quad p < 0.001 \quad s = 0.054 \\ SSY &= 0.688 \quad PRESS = 0.107 \quad R^2_{CV} = 0.844 \quad S_{PRESS} = 0.063 \quad S_{DEP} = 0.058 \end{aligned} \quad (5)$$

This Eqs. (4) and (5) explain 83.60% and 88.50% of the variances in the activity data. Significant values of R^2_{CV} (0.773 and 0.844 respectively) confirm the validity of these models. Lower values of S_{PRESS} suggest that optimum number of variables were taken to relate the antitumor activity (%TWI) of the structure of those analogs.

Table 7. t-statistic and p-values of QSAR Eqs. (1)–(7)

Eq.	Parameter/ Intercept	t-value	p-value	Eqn.	Parameter/ Intercept	t-value	p-value
1	Intercept	9.201	0.000	2	Intercept	9.653	0.000
	πR_4	5.404	0.000		πR_4	5.365	0.000
	$B_1 R_4$	3.784	0.001		$B_1 R_4$	3.724	0.001
			S_{AV}		-1.732	0.094	
3	Intercept	13.355	0.000	4	Intercept	12.020	0.000
	πR_4	3.556	0.001		<i>MRR</i> ₄	2.344	0.027
	$B_1 R_4$	6.749	0.000		$B_1 R_4$	6.032	0.000
	S_{AV}	-2.341	0.027		S_{AV}	-2.178	0.038
	<i>I</i>	-5.595	0.000		<i>I</i>	-5.639	0.000
5	Intercept	13.477	0.000	6	Intercept	13.283	0.000
	<i>LR</i> ₄	4.402	0.000		\mathcal{R}_1	2.570	0.016
	$B_1 R_4$	7.067	0.000		<i>LR</i> ₄	4.270	0.000
	S_{AV}	-2.631	0.014		$B_1 R_4$	7.099	0.000
	<i>I</i>	-6.651	0.000		<i>I</i>	-6.673	0.000
7	Intercept	13.096	0.000				
	\mathcal{R}_2	2.320	0.028				
	<i>LR</i> ₄	4.132	0.000				
	$B_1 R_4$	7.195	0.000				
	<i>I</i>	-6.612	0.000				

Another two models [Eqs. (6) and (7)] were developed using \mathcal{FR}_1 (field effect of the substituents at R_1 position) and \mathcal{RR}_2 (resonance effect of the substituent at R_2 position) instead of S_{AV} as shown below:

$$\begin{aligned} \text{Log (TWI)} &= 0.923 (\pm 0.069) + 0.127 (\pm 0.049) \mathcal{FR}_1 + 0.031 (\pm 0.007) LR_4 \\ &\quad + 0.335 (\pm 0.047) B_1R_4 - 0.173 (\pm 0.026) I \\ n = 32 \quad R^2 &= 0.884 \quad R^2_A = 0.866 \quad F_{(4,27)} = 51.495 \quad p < 0.001 \quad s = 0.054 \\ SSY &= 0.688 \quad PRESS = 0.093 \quad R^2_{CV} = 0.865 \quad S_{PRESS} = 0.059 \quad S_{DEP} = 0.054 \end{aligned} \quad (6)$$

$$\begin{aligned} \text{Log (TWI)} &= 0.958 (\pm 0.073) + 0.345 (\pm 0.149) \mathcal{RR}_2 + 0.031 (\pm 0.008) LR_4 \\ &\quad + 0.344 (\pm 0.048) B_1R_4 - 0.176 (\pm 0.027) I \\ n = 32 \quad R^2 &= 0.880 \quad R^2_A = 0.862 \quad F_{(4,27)} = 49.372 \quad p < 0.001 \quad s = 0.055 \\ SSY &= 0.688 \quad PRESS = 0.113 \quad R^2_{CV} = 0.840 \quad S_{PRESS} = 0.065 \quad S_{DEP} = 0.059 \end{aligned} \quad (7)$$

Eqs. (6) and (7) explain 88.40% and 88.00% variances of the activity data. Eq. (6) suggests that \mathcal{FR}_1 is conducive to the activity. Probably electron-donating group at R_1 position increases the field effect at this position. Eq. (7) indicates that \mathcal{RR}_2 is advantageous to the anticancer activity. These equations are also significantly predictive ($R^2_{CV} = 0.865$ and 0.840 respectively) in nature.

Table 8. Observed (Obs), Calculated (Calc) and LOO-predicted (Pred) activities of Eqs. (3)–(7)

Cpd	Obs	Eq. (3)		Eq. (4)		Eq. (5)		Eq. (6)		Eq. (7)	
		Calc	Pred	Calc	Pred	Calc	Pred	Calc	Pred	Calc	Pred
5	1.668	1.703	1.711	1.719	1.729	1.699	1.706	1.689	1.693	1.682	1.686
6	1.801	1.770	1.765	1.767	1.762	1.768	1.763	1.757	1.750	1.750	1.739
7	1.536	1.638	1.646	1.647	1.666	1.591	1.593	1.577	1.580	1.567	1.570
8	1.424	1.361	1.349	1.366	1.355	1.360	1.348	1.348	1.333	1.337	1.317
9	1.605	1.566	1.555	1.589	1.585	1.573	1.566	1.574	1.567	1.571	1.563
10	1.602	1.622	1.625	1.624	1.627	1.638	1.642	1.639	1.643	1.635	1.639
11	1.717	1.742	1.749	1.758	1.770	1.740	1.745	1.741	1.747	1.741	1.747
12	1.678	1.651	1.648	1.642	1.638	1.679	1.679	1.678	1.678	1.674	1.673
13	1.683	1.644	1.640	1.642	1.637	1.639	1.634	1.639	1.634	1.635	1.630
14	1.751	1.706	1.696	1.677	1.663	1.744	1.742	1.743	1.739	1.738	1.733
15	1.749	1.808	1.818	1.806	1.816	1.807	1.817	1.809	1.819	1.808	1.819
16	1.683	1.688	1.689	1.664	1.661	1.684	1.684	1.682	1.681	1.677	1.676
17	1.631	1.676	1.683	1.686	1.700	1.630	1.630	1.629	1.629	1.626	1.625
18	1.348	1.399	1.415	1.405	1.422	1.400	1.415	1.400	1.416	1.395	1.410
19	1.646	1.574	1.551	1.574	1.551	1.566	1.540	1.566	1.540	1.565	1.537
20	1.366	1.336	1.324	1.345	1.337	1.322	1.303	1.317	1.297	1.322	1.304
21	1.551	1.598	1.602	1.588	1.590	1.624	1.632	1.621	1.629	1.629	1.637
22	1.340	1.374	1.381	1.367	1.373	1.384	1.394	1.382	1.391	1.389	1.399
23	1.300	1.334	1.347	1.343	1.361	1.320	1.328	1.323	1.332	1.322	1.331
24	1.602	1.512	1.494	1.534	1.520	1.517	1.503	1.522	1.511	1.526	1.516
25	1.510	1.540	1.543	1.551	1.556	1.557	1.560	1.561	1.565	1.565	1.569
26	1.516	1.568	1.572	1.569	1.573	1.582	1.587	1.587	1.591	1.590	1.594
27	1.660	1.687	1.694	1.702	1.713	1.683	1.689	1.689	1.695	1.696	1.703
28	1.655	1.597	1.592	1.586	1.581	1.623	1.619	1.626	1.623	1.629	1.627
29	1.586	1.590	1.590	1.587	1.587	1.583	1.582	1.587	1.587	1.590	1.590
30	1.706	1.654	1.641	1.622	1.607	1.688	1.682	1.691	1.685	1.693	1.688
31	1.788	1.754	1.748	1.751	1.744	1.751	1.745	1.758	1.752	1.763	1.760
32	1.832	1.699	1.682	1.675	1.660	1.691	1.677	1.693	1.679	1.698	1.686
33	1.531	1.622	1.632	1.631	1.651	1.574	1.577	1.577	1.580	1.581	1.584
34	1.372	1.345	1.340	1.349	1.345	1.344	1.338	1.348	1.343	1.351	1.347
35	1.358	1.373	1.376	1.366	1.368	1.383	1.388	1.387	1.393	1.389	1.396
36	1.458	1.520	1.537	1.518	1.535	1.510	1.524	1.514	1.529	1.520	1.535

Confidence intervals of the final Eqs. (3)–(7) are more than 95% level as suggested by the *p*- and *t*-values shown in the Table 7. The observed, calculated and LOO-predicted activities of the Eqs. (3)–(7) are listed in the Table 8.

Table 9. Comparison among the Eqs. (3)–(7) and Eqs. (3a)–(7a)

	Eq. (3)	Eq. (3a)	Eq. (4)	Eq. (4a)	Eq. (5)	Eq. (5a)	Eq. (6)	Eq. (6a)	Eq. (7)	Eq. (7a)
<i>n</i>	32	26	32	26	32	26	32	26	32	26
<i>R</i> ²	0.865	0.877	0.836	0.854	0.885	0.914	0.884	0.916	0.880	0.912
<i>F</i>	43.512	37.605	34.443	30.794	52.041	56.321	51.495	57.170	49.372	54.092
<i>s</i>	0.059	0.054	0.065	0.059	0.054	0.045	0.054	0.045	0.055	0.046
<i>R</i> ² _{CV}	0.815	0.811	0.773	0.777	0.844	0.865	0.865	0.867	0.840	0.859
<i>S</i> _{DEP}	0.063	0.060	0.070	0.065	0.058	0.051	0.054	0.050	0.059	0.052
<i>R</i> ² _{Pred}	–	0.782	–	0.722	–	0.768	–	0.759	–	0.745

Table 10. Predicted activities of the compounds of the test set according to Eqs. (3a)–(7a)

Cpd	Observed activity	Predicted activity				
		Eq. (3a)	Eq. (4a)	Eq. (5a)	Eq. (6a)	Eq. (7a)
6	1.801	1.751	1.746	1.752	1.734	1.722
11	1.717	1.753	1.770	1.741	1.743	1.745
18	1.348	1.431	1.441	1.426	1.430	1.426
21	1.551	1.585	1.574	1.614	1.608	1.618
26	1.516	1.561	1.560	1.576	1.580	1.584
32	1.832	1.675	1.653	1.676	1.676	1.681

To prove the robustness of the QSAR models, validation was performed on a test set compounds by randomly excluding six compounds from the whole data set (*n* = 32) and regression was performed on the remaining compounds (*n* = 26) keeping the same combination of the independent parameters, *e.g.*, Eq. (3a) is the test set–training set model of Eq. (3) and so on. The excluded six compounds were then predicted using these new models [Eqs.(3a)–(7a)]. The comparison between these QSAR models is shown in Table 9. All of these test set–training set models are predictive in nature as shown in Table 9. Predictive activities of the test set compounds are listed in Table 10.

4 CONCLUSIONS

The QSAR study suggested that the length and width of the aliphatic substituent must be higher for the possible receptor ligand interaction through hydrophobic or dispersive force. The study also revealed that atoms 2, 3, 4, 5, 6, 7 are of great importance for anticancer activity of the glutamine analogs. Field effect of the R₁ substitutions possibly helps in electronic interaction with the substrate. In this case, electron-donating group may contribute in the binding with the electron deficient receptor site. Resonance effects of the R₂ substitution also might take part in the electronic interaction with the substrate. The presence of disubstitution at aliphatic side chain (R₄ and R₅) of glutamine analogs may not conducive to the activity. At least one free hydrogen atom in amide moiety may be essential for anticancer activity. This will help to select substituents for future synthesis of this type of compounds.

Acknowledgment

Authors are thankful to the University Grants Commission (UGC), New Delhi and All India Council for Technical Education (AICTE), New Delhi for providing financial assistance. Two of the authors BD and SS are grateful to the Council of Scientific and Industrial Research (CSIR), New Delhi and University Grants Commission (UGC), New Delhi for providing Senior Research Fellowship (SRF) and Junior Research Fellowship (JRF) respectively. Authors are also grateful to the authority of Jadavpur University for their help and encouragement.

5 REFERENCES

- [1] M. Medina, Glutamine and cancer, *J. Nutr.* **2001**, *131*, 2539S–2542S.
- [2] M. A. Medina, F. Sánchez-Jiménez, J. Márquez, A. R. Quesada and I. Núñez de Castro, Relevance of glutamine metabolism to tumor cell growth, *Mol. Cell. Biochem.* **1992**, *113*, 1–15.
- [3] W. W. Souba, Glutamine and cancer, *Ann. Surg.* **1993**, *218*, 715–728.
- [4] V. S. Klimberg and J. L. McClellan, Glutamine, cancer, and its therapy, *Am. J. Surg.* **1996**, *172*, 418–424.
- [5] D. L. Bartlett and S. Charland, Effect of glutamine on tumor and host growth, *Ann. Surg. Oncol.* **1995**, *2*, 71–76.
- [6] T. R. Austgen, P. S. Dudrick and H. Sitren, The effects of glutamine-enriched total parenteral nutrition on tumor growth and host tissues, *Ann. Surg.* **1992**, *215*, 107–113.
- [7] C. Costa, J.F. Huneau and D. Tome, Characteristics of L-glutamine transport during Caco-2 cell differentiation, *Biochem. Biophys. Acta/Biomemb.* **2000**, *1509*, 95–102.
- [8] C. L. Collins, M. Wasa, W. W. Sauba, and S. F. Abcouwer, Determinants of glutamine dependence and utilization by normal and tumor-derived breast cell lines, *J. Cell. Physiol.* **1998**, *176*, 166–178.
- [9] S. Sukemori and K. Sugimura, Inhibition of cultured leukemia cell growth by enhanced adriamycin cytotoxicity with reduction of glutamine or asparagine level in medium, *Cancer Biochem. Biophys.* **1994**, *14*, 99–105.
- [10] F. Devreker, R. M. Winston and K. Hardy, Glutamine improves human preimplantation development in vitro, *Fertil. Steril.* **1998**, *69*, 293–299.
- [11] G. Ollenschlager, A. Simmel and E. Roth, Availability of glutamine from peptides and acetylglutamine for human tumor-cell cultures, *Metabolism*, **1989**, *38*, 40–42.
- [12] K. Srikanth, C. A. Kumar, D. Goswami, A. U. De and T. Jha, Quantitative structure activity relationship (QSAR) studies of some substituted benzenesulphonyl glutamines as tumour suppressors, *Ind. J. Biochem. Biophys.* **2001**, *38*, 120–123.
- [13] K. Srikanth, B. Debnath and T. Jha, Syntheses, biological evaluation and QSAR study on antitumor activity of 1,5-*N,N'*-disubstituted-2-(substituted benzenesulphonyl) glutamamides, *Bioorg. Med. Chem.* **2002**, *10*, 1841–1854.
- [14] K. Srikanth, C. A. Kumar, B. Ghosh and T. Jha, Synthesis, screening and quantitative structure-activity relationship (QSAR) studies of some glutamine analogues for possible anticancer activity, *Bioorg. Med. Chem.* **2002**, *10*, 2119–2131.
- [15] B. Debnath, K. Srikanth, S. Banerjee and T. Jha, 1,5-*N,N'*-Disubstituted-2-(substituted benzenesulphonyl)-glutamamides as antitumor agents. part 2. synthesis, biological activity and QSAR study, *Internet Electron. J. Mol. Des.* **2002**, *1*, 488–502. www.biochempress.com
- [16] T. Jha, B. Debnath, S. Samanta and A. U. De, QSAR study on some substituted glutamine analogs as anticancer agents, *Internet Electron. J. Mol. Des.* **2003**, *2*, 539–545. www.biochempress.com
- [17] S. Samanta, K. Srikanth, S. Banarjee, B. Debnath, S. Gayen and T. Jha, 5-*N*-Substituted-2-(substituted benzenesulphonyl) glutamines as antitumor agents II: synthesis, biological activity and QSAR study, *Bioorg. Med. Chem.* **2004**, *12*, 1413–1423.
- [18] F. Hannaford and S. Tatchell, In: Vogel's Textbook of Practical Organic Chemistry; ELBS/Longman: Singapore, 1996; p877.
- [19] J. March, (Ed). In Advanced Organic Chemistry, Reactions, Mechanism and Structure; John Wiley & Sons: New York, 1992; p 417.
- [20] S. K. Purkayastha, T. Jha, D. K. Pal and A. U. De, Possible antineoplastic agents: Part XIII. Synthesis, biological evaluation and QSAR studies of some 1-(substituted benzenesulphonyl)-5-oxopyrrolidine-2-carboxylic acid derivatives, *Anticancer Drug Des.* **1993**, *8*, 95–100.
- [21] H. Kubinyi, QSAR: Hansch analysis and related approaches, In: R. Mannhold, P. Krogaard-Larsen and H. Timmerman, (Ed.), Methods and Principles in Medicinal Chemistry, **1993**, VCH, New York.
- [22] C. Hansch and A. Leo, *Substituent constants for correlation analysis in chemistry and biology*, **1979**, John Wiley & Sons: NY, pp 48–63.
- [23] L. B. Kier and L. H. Hall, An electrotopological-state index for atoms in molecules, *Pharm. Res.* **1990**, *7*, 801–

807.

- [24] K. Rose, L. H. Hall and L. B. Kier, Modeling blood–brain barrier partitioning using the electrotopological state, *J. Chem. Inf. Comput. Sci.* **2002**, *42*, 651–666.
- [25] Mouse, a computer programme written in C⁺⁺ language by S. K. Naskar of Department of Computer Science and Engineering, Jadavpur University and B. Debnath, one of the authors of this article.
- [26] G. W. Snedecor and W.G. Cochran, *Statistical Methods*; Oxford & IBH: New Delhi, 1967, pp. 381.
- [27] J. O. Rawlings, *Applied Regression Analysis*; Wardsworth: CA, USA, 1988, pp. 327.
- [28] N. R. Darper and H. Smith, *Applied Regression Analysis*, John Wiley & Sons: New York, 1981, pp. 294.
- [29] E. J. Lien, In *Quantitative Correlations Using Multiple Regression and Substituent Constants*; E. J. Ariens, Drug Design; Academic: New York, 1975; Vol. V, pp. 84.
- [30] Multiregress, a computer programme written in C⁺⁺ language by S. K. Naskar of Department of Computer Science and Engineering, Jadavpur University and B. Debnath, one of the authors of this article.
- [31] K. Srikanth, B. Debnath and T. Jha, QSAR study on adenosine kinase inhibition of pyrrolo[2,3-*d*] pyrimidine nucleoside analogs using Hansch approach, *Bioorg. Med. Chem. Lett.* **2002**, *12*, 899–902.
- [32] B. Debnath, S. P. Vishnoi, B. Sa and T. Jha, QSAR study on some dihydrofolate reductase inhibitors, *Internet Electron. J. Mol. Des.* **2003**, *1*, 128–136. www.biochempress.com
- [33] B. Debnath, S. Samanta, K. Roy and T. Jha, QSAR study on some *p*-arythio cinnamides as antagonists of biochemical ICAM–1/LFA–1 interaction and ICAM–1/JY–8 cell adhesion in relation to anti–inflammatory activity, *Bioorg. Med. Chem.* **2003**, *11*, 1615–1619.
- [34] B. Debnath, S. Samanta, S. K. Naskar, K. Roy and T. Jha, study on the affinity of some arylpiperazines towards the 5-HT_{1A}/α₁–adrenergic receptor using the *E*–state index, *Bioorg. Med. Chem. Lett.* **2003**, *13*, 2837–2842.
- [35] B. Debnath, S. Gayen, S. K. Naskar, K. Roy and T. Jha, QSAR study of some azidopyridinyl neonicotinoids insecticides for their selective affinity towards the *Drosophilla* nicotinic receptor over mammalian α₄β₂ receptor using electrotopological state atom index, *Drug Des. Discov.* **2003**, *18*, 81–89.
- [36] B. Debnath, S. Gayen, S. Bhattacharya, S. Samanta and T. Jha, QSAR study on some pyridoacridine ascididemin analogs as anti–tumor agents, *Bioorg. Med. Chem.* **2003**, *11*, 5493–5499.
- [37] S. Gayen, B. Debnath, S. Samanta and T. Jha, QSAR study on some anti–HIV HEPT analogues using physicochemical and topological parameters, *Bioorg. Med. Chem.* **2004**, *12*, 1493–1503.
- [38] B. Debnath, S. Gayen, A. Basu, K. Srikanth and T. Jha, Quantitative structure–activity relationship study on some benzodiazepine derivatives as anti–Alzheimer agents. *J. Mol. Mod.* **2004**, in press.
- [39] B. Debnath, S. Gayen, A. Basu, B. Ghosh, K. Srikanth and T. Jha, Quantitative Structure–Activity Relationship Study Using Refractotopological State Atom Index on Some Neonicotinoid Insecticides, *Bioorg. Med. Chem.* **2004**, *12*, 6137–6145.